

Studies on composition and components of air spores belonging to phycomycetes over the field of sugarcane

Kadam SS¹, Kabadi S N²

¹Department of Botany, ²Department of Chemistry, ACS College, Shankarnagar, Dist. Nanded, Maharashtra, INDIA.
Email: kadamssunil2371@gmail.com, sambhaji456@gmail.com

Abstract During the present studies composition and components of the air spores belonging to the Phycomycetes over the field of sugarcane. Slides of spore catches were prepared, mounted and spore scanning was carried out regularly. Four types of fungal spores belonging to the Phycomycetes were trapped on the cellophane tape fixed on the drum of the sampler. Apart from these various dust particles were also seen in abundance.

Key Words: Air spores, phycomycetes, sugarcane, fungal spores, and cellophane.

*Address for Correspondence:

Dr. Kadam SS, Department of Botany, ACS College, Shankarnagar, Dist. Nanded, Maharashtra, INDIA.
Email: kadamssunil2371@gmail.com

Access this article online

Quick Response Code:



Website:

www.statperson.com

Accessed Date:

10 March 2018

INTRODUCTION

Aerobiology is an interdisciplinary science which deals with the study of biological components like pollen grains, fragments of fungal spores, hyphal fragments, bacteria, viruses, algae, lichens, plant seeds and other propagules, protozoa, minute insects and insect parts etc. in the atmosphere. A biotic particulate and gas affecting living organisms have been recently included in the concept of aerobiology. The aerobiological studies are mainly concerned with the interrelationships between the biological components in the atmosphere, sources of biological components, their release in the atmosphere, their deposition and impact on health of plants and animals including human beings. Airborne infections and the resulting diseases threaten the lives and productivity of plants. Airborne diseases still pose a challenge to mankind. It is evident from the literature that investigations on outdoor (extramural) atmosphere for the

detection of microorganisms have got manifold significance and utility. The aero

allergenic pollen grains, fungal spores and other biologically significant particles detected from the atmosphere have their impact on

human health. The detection and trapping of these significant plant pathogenic airborne fungi over crop fields are ultimately helpful in formulating a better and efficient forecasting system. It is also clear from the literature that many workers have focused attention towards the relationship between airborne fungi and phytopathological problems. In India several workers like Cunningham (1873), Mehta (1952), Padmanabhan (1953), Ramalingam (1966), Sreeramulu (1970), and Tilak (1970) carried out systematic and classical aerobiological work on rust spores with reference to wheat, Jowar, bajra, rice, sugarcane, cotton, banana, citrus and vegetables at various research centers.

In Maharashtra particularly in the Marathwada region aerobiological work with reference to many crop plants was extensively carried out by various workers like Tilak (1970), Nagpure (1973), Bhalke (1979), Chakre (1979), Balkhanale (1980), Babu (1983), Meshram (1990), Naik (1995), Pawar (1997), Kadam (2002), Bembrekar (2005), Dhaware (2007) and Patil (2008) at various centres. However, detailed studies on air spores over Botanical gardens in respect of seasonal conditions are meager. Considering these facts studies on air spores belonging to the Phycomycetes over the field of sugarcane is carried out.

MATERIALS AND METHODS

Composition of air spores belonging to Phycomycetes over the field of sugarcane during two consecutive kharif seasons (Table-1, Plate-

2): In order to study composition and components of the air spores belonging to the Phycomycetes over the botanical garden, slide of spore catches were prepared, mounted and spores scanning was carried out regularly. Before the scanning, the slides were marked with a ball-

point pen into six equal parts, each part indicating the spore catch of two hours of sampling period. Area of 9600 sq. micron of the total area of the trace obtained was scanned under 10X x 45X eyepiece objective combination of binocular research microscope. The identification of fungal spore types was made on the basis of morphological characters, visual identification by comparison with reference slides.

RESULTS AND DISCUSSION

Analysis of spore catches from the results presented in table-1 revealed that four types of fungal spores belonging to the phycomycetes were trapped on the cellophane tape fixed on the drum of the sampler. Apart from these various dust particles were also seen in abundance. Enumeration of the identified spores of fungi belonging to Phycomycetes over the Sugarcane field during the two consecutive kharif seasons:

Acaulopage Drechs: Mycelium slender, aseptate, sparsely branched. Conidia aerial, hyaline, borne singly on prostrate hyphae. During the present studies the spore concentration in the first test kharif season from 21-06-2009 to 30-09-

2009 and second test kharif season from 21-06-2010 to 30-09-2010 was recorded 1498/m³ and 1190/m³ of air respectively. They contributed 0.91% and 0.74% to the total air spore in the first and second kharif seasons respectively. The average percentage contribution for both the test seasons was recorded 0.82% to the total air spore over the botanical garden.

Albugo Pers: The spore one celled, nearly globose, thick walled, smooth, hyaline to yellow coloured, 14-

16 µm. borne in chain on aseptate club shaped conidiophores. During the present studies the spore concentration in the first test kharif season from 21-06-2009 to 30-09-2009 and second test kharif season from 21-06-2010 to 30-09-2010 was recorded 4830/m³ and 5334/m³ of air respectively. They contributed 2.95% and 3.32% to the total air spore in the first and second kharif seasons respectively. The average percentage contribution for both the test seasons was recorded 4.61% to the total air spore over the Sugarcane field.

Mucor Micheliex Fr: Spore one celled globose to oval, thin walled, smooth dark in mass 3-

5 µm diameter. common during wet months of rainy seasons. saprophytic on a variety of substrates. During the present studies the

spore concentration in the first test kharif season from 21-06-2009 to 30-09-2009 and second test kharif season from 21-06-2010 to 30-09-

2010 was recorded 2408/m³ and 238/m³ of air respectively. They contributed 1.47% and 0.14% to the total air spore in the first and second kharif seasons respectively. The average percentage contribution for both the test seasons was recorded 0.80% to the total air spore over the Sugarcane field.

Rhizopus Ehren: The spores of *Rhizopus* are unequal, irregular, round, oval, angular, striate, 9-

11 x 7.5 µm in size. It is common saprophyte and facultative parasite on mature fruits, vegetables and on many other substrates. During the present studies the spore concentration in the first test kharif season from 21-06-2009 to 30-09-

2009 and second test kharif season from 21-06-2010 to 30-09-2010 was recorded 462/m³ and 2772/m³ of air respectively. They contributed 0.28% and 1.72% to the total air spore in the first and second kharif seasons respectively. The average percentage contribution for both the test seasons was recorded 1% to the total air spore over the Sugarcane field.

Table 1: Composition of air spores belonging to Phycomycetes over the Sugarcane field during two consecutive kharif seasons

Sr.No.	Spore type
1	<i>Acaulopage Drechs</i>
2	<i>Albugo Pers.</i>
3	<i>Mucor Micheliex Fr.</i>
4	<i>Rhizopus Ehren.</i>




Figure 1:

REFERENCES

1. Babu, M. (1983): Aerobiological studies at Aurangabad. Ph.D. Thesis, Marathwada University, Aurangabad.
2. Bembrekar, S.K. (2005): Studies in air spore of cotton field. Ph.D. Thesis, Swami Ramanand Teerth Marathwada University, Nanded.

3. Bhalke, S.P. (1981): Airspora over some fields. Ph.D. Thesis, Marathwada University, Aurangabad.
4. Chakre, O.J. (1979): Studies in microbial population of fair at Aurangabad. Ph.D. Thesis, Marathwada University, Aurangabad.
5. Cunningham, D.D. (1873): Microscopic examination of fair. Govt. Printer, Dhaware, D.A. (2007): study of aeromycoflora and Phyllosphere flora over some edible fruit plants, Ph.D. Thesis, Swami Ramanand Teerth Marathwada University, Nanded. Calcutta, pp 58.
6. Kadam, S.S. (2002): Aerobiological investigation over sugarcane fields at Shankarnagar.
7. Mehta, K.C. (1952): Further studies of cereal rusts in India-II. Sci. Monogr. Coun. Agric. Res. India 18: 1-368.
8. Meshram, B.M. (1990): Studies in airspora at Udgir. Ph.D. Thesis, Marathwada University, Aurangabad.
9. Nagpurne, S.D. (1973): Studies in airspora at Kandhar. Ph.D. Thesis, Marathwada University, Aurangabad.
10. Naik, M.N. (1995): Atmospheric microbial population studies at Aurangabad (M.S.). Ph.D. Thesis, Marathwada University, Aurangabad.
11. Padmanabhan, S.Y. (1953): Specialization in pathogenicity of *Helminthosporium oryzae*. Proc. 40th Indian Sci. Congr. III: 18.
12. Patil, T.M. (2008): Studies in airspora over black gram fields. Ph.D. Thesis, Dr. Babasaheb Ambedkar Marathwada University, Aurangabad.
13. Pawar Dr. Babasaheb Ambedkar Marathwada University, Aurangabad.
14. Ramalingam, A.A. (1966): A volumetric survey of the atmosphereic pollen over paddy fields at Visakhapat, S.R. (1997): Studies in airspora over some crop fields at Aurangabad. Ph.D. Thesis, namin 1960 and 1961. Palynol. Bull. 2 and 3: 11-17.
15. Sreeramulu, T. (1970): Conidial dispersal in two species of *Cercospora* causing Tikka leaf spots on groundnut (*Arachis hypogaea*). J. Agri. Sci. 40: 173-178.
16. Tilak, S.T. (1984): Aerobiology and cereal crop diseases. Rev. Trop. Pl. Path. (Ed. S.P. Raychaudhari) 1: 329-354.
17. Tilak, S.T. (1989): Aeropalynology of Maharashtra. Recent Advances in Palynology. 101-116.
18. Tilak, S.T. (1989): Airborne airborn pollen and fungal spores. Vijayanti prakashan, Aurangabad.
19. Tilak, S.T. (1991): Fungal spora and allergy. J. Palynol. 27. p. 36-386.
20. Tilak, S.T. and M. Babu (1982): Incidence of *Cunninghamella* spores in the atmosphere. Poll. Res., 1: 55-58.
21. Tilak, S.T. and S.P. Bhalke (1978): Aeromycology at Aurangabad-II Deuteromycetes. Indian J. Bot., 1: 113-115.
22. Tilak, S.T. and B.V. Srinivasulu (1967): Airspora of Aurangabad. Ind. J. Microbiology; 7: 167-170.
23. Tilak, S.T. and B.V. Srinivasulu (1967): Airspora of Aurangabad. Indian J. Microbiol. 7: 167-170.
24. Tilak, S.T. and M. Babu (1982): Incidence of *Cunninghamella* spores in the atmosphere. Poll. Res. 1: 55-58. Tilak and S.P. Bhalke (1978): Aeromycology at Aurangabad.-II. Deuteromycetes. Indian J. Bot. 1: 113-115.
25. Tilak, S.T. and O.J. Chakre (1978): Atmospheric concentration of *Claviceps microcephala* over bajra fields in relation to their environmental factors. In Int. Palyno. con. Lucknow. (Abstract.) 181-182.
26. Tilak, S.T. and R. Kulkarni (1970): A new air sampler. Experimentia, 26: 443.
27. Tilak, S.T. and R.L. Kulkarni (1970): A new air sampler. Experimental. 26: 443.
28. Tilak, S.T. and R.L. Kulkarni (1971): Airspora over sugarcane fields. International Symp. Patho. New Delhi.
29. Tilak, S.T. and R.L. Kulkarni (1975): Atmospheric concentration of *Rhizopus* spores in air. Biovigyanam. 1: 91-94.
30. Tilak, S.T. and K.S. Ramchander Rao (1989): Aerobiology of S. unflower disease. Environmental Ecology and Aerobiology. edited by S.T. Tilak, Vol. 3: 41-66.
31. Tilak, S.T. (1980): Airborne pollen and fungal spores. Vijayanti Prakashan, Aurangabad.
32. Tilak, S.T. (1982): Aerobiology and its practical applications. Proc. INat. Conf. Env. Bio. 33-38.
33. Tilak, S.T. and B.V. Srinivasulu (1967): Airspora of Aurangabad. Indian J. Microbiol. 7: 167-170.
34. Tilak, S.T. and D.B. Vishwe (1975): microbial content of air in a delibary. Biovigyanam. 1: 91-94.
35. Tilak, S.T. and S.P. Bhalke (1981): Aeromycology at Aurangabad-III Hyphal fragments. Advancing Frontiers of Mycol. and Pathol. Today and Tomorrow Publ. Delhi, pp. 55-51.
36. Tilak, S.T. (1987): Atlas of airborn pollen and fungal spores. Vijayanti prakashan, Aurangabad.
37. Tilak, S.T. and K.S. Ramchander Rao (1989): Trapping *Cunninghamella* spores from air. Poll. Res. 18(1): 37-40.

Source of Support: None Declared
Conflict of Interest: None Declared